

**PRESETTLEMENT BARRENS IN THE
GLACIATED PRAIRIE REGION OF ILLINOIS¹**

Marlin L. Bowles & Jeanette L. McBride

*Proceedings
of the*

**NORTH AMERICAN
CONFERENCE ON
SAVANNAS AND BARRENS**



*Illinois State University
Normal, Illinois
October 15-16, 1994*

PRESETTLEMENT BARRENS IN THE GLACIATED PRAIRIE REGION OF ILLINOIS¹

Marlin L. Bowles & Jeanette L. McBride

"About half a mile from the cabin we passed a small prairie, and soon entered the woods again, and then some barrens. A barren island nearly destitute of timber, but overrun with scrubby underwood. After passing these barrens for more than a mile, we got sight of the prairies" (Woods 1822).

INTRODUCTION

Classification, management and restoration of natural plant communities are difficult objectives for Midwestern conservation agencies, especially when the structure and composition of former vegetation are unclear and management decisions are affected by nomenclature. For example, **prairie**, **savanna**, and **timber** are thought to have comprised the presettlement vegetation mosaic in the forest-prairie transition of Illinois, with savanna occurring in areas of intermediate fire protection (Kilburn 1959, Moran 1976, 1980, Anderson 1991, Bowles *et al.* 1994). Because little specific information is available about presettlement oak savanna and woodland (Apfelbaum & Haney 1991), they have become the focus of community classification (Faber-Langendoen *in press*), conservation and restoration (*e.g.* Burger *et al.* 1991, Stearns *in press*), and research on groundlayer vegetation (Packard 1991, Leach 1994).

This paper treats the related but vaguely classified and poorly understood **barrens** plant community. Although often considered a feature of unglaciated regions, barrens were frequently referred to in the glaciated Illinois landscape at the time of settlement (Vestal 1936, Hutchison 1987). The Illinois Natural Areas Inventory (INAI) classified barrens as a subtype of savanna usually caused by topo-edaphic features such as thin soils over bedrock, and occurring as local inclusions of prairie flora in forested southern and western Illinois (White & Madany 1978, Madany 1981, Anderson & Schwegman 1991). Madany (1981) had 24% floristic similarity between savannas and barrens (exclusive of sand savanna) sampled by the INAI, while Betz (1992) found less than 30% similarity between cemetery savanna floras and the "barrens" flora of Mead (1846). However, Packard (1991) contended that barrens as observed by Mead (1846) were synonymous with savanna.

OBJECTIVES

Do glaciated barrens fall within a savanna concept of non-overlapping canopy tree crowns with a graminoid fuel matrix (*e.g.* Stout 1946, Bray 1958, Curtis 1959, White & Madany 1978, Madany 1981, Packard 1993), or do they represent a different plant community? Is more specific information available about the former structure and ecology of barrens in the prairie region of Illinois? To address these questions, we review historic descriptions and species, classify barrens in glaciated Illinois, and present a model for barrens

¹1994. Pages 75-85 *in* Proceedings of the North American Conference on Savannas and Barrens (J.S. Fralish, R.C. Anderson, J.E. Ebinger, & R. Szafoni, eds.), Illinois State University, Normal.

development within a prairie-forest continuum. This paper excludes sand substrates and focuses on silt loam soils, which comprised the substrate for Illinois tallgrass prairie.

HISTORIC DESCRIPTIONS

Terminology

"Barrens" is one of the more ambiguous eastern North American vegetation terms, but usually connotes stressful habitats (*e.g.* White & Madany 1978, Homoya 1994). It has described vegetation differently across regions and over time. For example, the term has been used to classify natural pine and oak sand barrens in Wisconsin (Stout 1946, Curtis 1959), naturally burned heathlands such as the Pine Barrens of New Jersey (Matlack *et al.* 1993), natural or anthropogenic barrens caused by substrate, logging, and fire in Nova Scotia (Hall & Aalders 1968, Strang 1972), and forest openings caused by fire or edaphic factors in the southeastern U.S. (*e.g.* DeSelm 1981, 1986; Homoya 1994). Schroeder (1982) interpreted barrens as an Appalachian term applied to sites perceived as unproductive because of failure to grow trees. Williams (1981) generalized that barrens was applied historically to shrub savannas located chiefly south of the northeasterly axis of the prairie peninsula, while "oak openings," or tree savannas, occurred primarily to the north. Indeed, shrubs and oak grubs occurred in association with grassland and forest openings in sites described as barrens in southern Illinois (*e.g.* Engelman 1863, Hutchison *et al.* 1986), Indiana (Deam 1940, Keith 1980) and in Kentucky (*reviewed in* Baskin & Baskin 1981).

As settlement and the Public Land Survey entered the extensive glaciated prairies of Illinois, barrens terminology was applied to perceived unproductive habitats with vegetation intermediate between grassland and forest, often with "tall grass," but characterized by "underwood" or "brushwood" of fire-resistant shrubs or oak grubs (*e.g.* Birkbeck 1818, Woods 1822, Peck 1834), rather than the open-grown trees of savanna oak openings (*sensu* Gleason 1922, Stout 1946, Curtis 1959, Bray 1960). Barrens in St. Louis (Beck 1826, 1828) were "undulating," with "shrubby oak" and hazel "shrubbery" (*reviewed in* Schroeder 1982). In Illinois, Mitchell (1838) described "bushy" prairies in Illinois covered with hazel and furze, small sassafras, shrubs, and grape vines. Short (1845) referred to sparse oak cover with dense undergrowth of shrubs in barrens, which he used synonymously with hazel "roughs" and "thickets." Farnham (1846) referred to "copse" growth of hazel in barrens, and Brendel (1887) listed vegetation of copses. Bebb (1860) also referred to the hazel undergrowth extending a mile or more into the prairie forming a "hazel ruff."

Ecological factors: fire, topography, and successional change

Although Hutchison (1986) and Homoya (1994) contend that barrens generally have edaphically restricted plant growth, reduced herbaceous diversity, and ecological stability, most historic accounts (*reviewed in* White 1994) describe barrens as rich and productive, and maintained by fire (*e.g.* Peck 1834, Mitchell 1838). According to Gleason (1922), Illinois barrens represented a late stage of fire-caused forest degeneration characterized by four- to

five- foot high sprouts of scrub oak (apparently *Quercus velutina*), hazel (*Corylus americana*), and wild plum (*Prunus americana*). They were most common in uneven or rolling topography or stream drainages (e.g. Peck 1834), which reduced fire effects, or developed on the west sides of forests attacked by eastward-moving prairie fires driven by prevailing winds (Gleason 1913). Thus, north or east sides of prairie groves may have had more stable forest-prairie margins.

With advancing settlement and fire protection, many authors described successional instability of barrens (White 1994). In a remarkably lucid description, Peck (1834) noted that "when the fires are stopped, these barrens produce timber ...", from a "...mass of roots ...," and that when fire protection and grazing reduced prairie grasses, the prairies "begin to assume the character of barrens; first hazel and other shrubs, and finally a thicket of young timber..." By 1873, Ridgway (1873) remarked that "prairies of often ten miles or more .. are now entirely overgrown with a dense scrub of hazel, ..." and that "twenty years from now, they will have ... become transformed into the usual woods of the region." Thus, large areas of barrens were converted into forest "as by magic" when the anthropogenic fires that had maintained them were stopped and the oak sprouts became trees (Gleason 1922).

Species composition: shrub and oak grub dominance

Shrubs and fire-stunted oak grubs appear to have been structurally dominant components of barrens. Historic descriptions (Woods 1822, Peck 1834, Mitchell 1839, Short 1845, Gerhard 1857, Bebb 1860, Hilgard *n.d.*), identify more than 30 shrub species that may have characterized barrens, including hazel, New Jersey tea (*Ceanothus americanus*), the dogwoods *Cornus drummondii*, *C. obliqua*, *C. racemosa*, & *C. stolonifera*, wild crab (*Malus coronaria* & *M. ioensis*) wild plum (*Prunus americana*), sumac (*Rhus* sp.), rose (*Rosa* sp.), prairie willow (*Salix humilis*), and prickly ash (*Zanthoxylum americanum*). Barrens formed along the western flanks of forests were dominated by hazel, with *Quercus velutina* and *Q. imbricaria* forming the interior forest margin (Gleason 1913).

As with shrubs, historic descriptions list more than 30 forb species occurring in barrens. If these lists focus on showy or distinctive species, they may not represent a typical flora. To obtain a more comprehensive historic list of species that occurred in barrens, we analyzed four historic annotated floras that contained references to either barrens (Mead 1846), "copse" or "thicket" (Higley & Raddin 1871), "copse" (Brendel 1887), or "thicket" (Hus 1908), which appear to be analogous or similar habitats. These floras include counties in which barrens were recorded by the Public Land Survey (Hutchison 1987, M.L. Bowles, pers. obs.). The resulting list (Appendix I) contains 243 taxa that apparently occurred in glaciated barrens. Herbaceous and graminoid species include *Asclepias purpurascens*, *Baptisia leucantha*, seven *Desmodium* species, *Echinacea purpurea*, four *Eupatorium* species, seven *Helianthus* species, four *Lespedeza* species, two *Elymus* species, five *Panicum* species, seven *Solidago* species, three *Pycnanthemum* species, and the blazing star *Liatris scariosa* var. *nieuwlandii*. The abundance and diversity of grass species is unclear. Although some prairie and woodland grasses are listed, nine of ten grass species that Packard (1991) proposed for addition to the Mead's barrens flora are absent. Even though "tall grass"

² Nomenclature follows: Jones, G.N. & G.D. Fuller. 1955. Vascular plants of Illinois. University of Illinois Press, Urbana & Illinois State Museum, Springfield (Scientific Series, Vol VI).

is described for barrens, Hilgard (*n.d.*) noted that "within these shrubby areas the grass was much suppressed." Gleason (1913) assumed that grasses persisted within hazel zones, but admitted that "there seem to be no authentic data on the matter."

A CLASSIFICATION AND MODEL FOR GLACIATED PRAIRIE BARRENS

Classification

Historic descriptions of barrens on glaciated soils in the prairie region of Illinois are analogous to two related and intergrading community types that were only partially classified by the INAI or Curtis (1959). One type, **shrub prairie**, was defined by the INAI as comprising hazel and plum thickets on silt soils, but no such natural remnants were found (White & Madany 1978). The second type, **oak grub** or **oak brush savanna** (Curtis 1959), had oaks reduced to grub sprouts by repeated fires. On silt soils, the INAI recognized savanna only with mature trees forming a partial canopy cover. We suggest an alternative classification that would combine the two types into a **prairie barrens** category with **shrub barrens** and **oak grub barrens** subtypes that commonly merge into a single community.

Conceptual model

Figure 1 presents a conceptual landscape model for development of barrens and savanna on glaciated silt soils. Although these communities probably co-occurred or integrated at a local scale, this model provides separate pathways for understanding their development. The different factors responsible for shrub barrens, oak grub barrens, and savanna are not clear. References by the Public Land Survey to timber "with" or "without" undergrowth in different landscape positions (Bowles *et al.* 1994) suggest that large scale ecological processes were important. According to Gleason (1922), the parklike conditions known as oak openings (savanna) resulted from destruction of forest undergrowth and susceptible species by fire, and invasion by prairie. This is modeled as a "Savanna Pathway," by which woody understory but not overstory is lost to fire. The fire-resistant bur oak (*Quercus macrocarpa*) would be most adaptable to this process, while other oaks that are less fire resistant but sprout after fire would be reduced to oak grub barrens (Curtis 1959). Development of a "Barrens Pathway," by which woody understory vegetation persists while canopy trees are lost is less clear and the exact cause was unknown to Gleason (1922). Sprouting of fire-adapted hazel and oaks, and absence of fire resistant bur oak may have contributed to this process, while immigration of animal- or wind-dispersed woody species such as plumb, crabs, or willow would have added species to barrens (Gleason 1922). Landscape fire effects probably contributed to these different pathways. As indicated, barrens were usually associated with more rugged topography, which formed partial fire breaks. For example, Bowles *et al.* (1994: Figure 1) found presettlement barrens associated with an area of the West Chicago Moraine in DuPage Co., IL that was not fire protected by the DuPage River. Spatial dynamics in this model are driven by temporal climatic variation. Barrens and savanna encroach toward forest as warm dry conditions promote fire, and toward prairie with cool wet conditions that promote fire protection.

CONCLUSIONS

Presettlement barrens on silt soils in glaciated Illinois were apparently misconstrued as sterile or unproductive, as their vegetation quickly transformed to forest with fire protection. Barrens and savanna differed by presence or absence of woody understory vegetation, and formed through forest degeneration under conceptually different ecological pathways affected by landscape features, fire, and species characteristics. Barrens appear to have been shrub- and oak grub-dominated with limited or patchy distribution of grasses, while a stronger component of prairie vegetation and grasses occurred in savanna. Based on historic descriptions, it appears that barrens were spatially and temporally dynamic. They rapidly succeeded to forest with fire protection, but also manifested in prairie from shrub or grub sprouts that were suppressed by anthropogenic fire. Without fire protection or Indian-set fires, climatic variation would have caused temporal shifts in the spatial pattern of this vegetation. Misapplication of savanna terminology to barrens natural areas could result in management to remove native shrub understories that belong in these communities.

ACKNOWLEDGMENTS

We thank Robert Betz, Don Faber-Langendoen, and Jack White for sharing ideas on the concept of glaciated barrens, and Mary K. Solecki, Jack White, Jerry Baskin, Mike Homoya, and Christopher Dunn for comments on various parts of this manuscript.

LITERATURE CITED

- Anderson, R.C. & J.E. Schwegman. 1991. Twenty years of vegetational change on a southern Illinois barren. *Natural Areas Journal*. 11:100-107.
- Anderson, R.C. 1991. Presettlement forests of Illinois. Pages 9-19 in: Proceedings of the Oak Woods Management. G.V. Burger, J.E. Ebinger, and G.S. Wilhelm, eds. Workshop. Eastern Illinois University, Charleston.
- Apfelbaum, S.I. & A. W. Haney. 1991. Management of degraded oak savanna remnants in the upper Midwest: preliminary results from three years of study. Pages 81-90 in: Proceedings of the Oak Woods Management Workshop. G.V. Burger, J.E. Ebinger, & G.S. Wilhelm, eds. Eastern Illinois University, Charleston.
- Baskin, J.M. & C.C. Baskin. 1981. the big barrens of Kentucky not a part of Transeau's prairie peninsula, Pages 43-48 in: Proceedings of the Sixth North American Prairie Conference. Ronald Stuckey & K.J. Reese, eds. Ohio Biological Survey Notes No. 15.
- Bebb, M.S. 1860. The flora of Ogle and Winnebago Counties., Ill. *The Prairie Farmer*. 22(6):182-183.
- Beck L.C. 1826. Contributions towards the botany of the states of Illinois and Missouri. *American Journal of Science and Arts* 10:257-264).
- Beck L.C. 1828. Contributions towards the botany of the states of Illinois and Missouri. *American Journal of Science and Arts* 11:167-182.
- Betz, R.F. 1992. Species composition of old settler savanna and sand prairie cemeteries in northern Illinois and Northwestern Indiana. Pages 79-87 in: Proceedings of the

- Twelfth North American Prairie Conference. D.D. Smith & C.A. Jacobs, eds. University of Northern Iowa, Cedar Falls.
- Birkbeck, M. 1818, Notes on a journey in America from the coast of Virginia to the territory of Illinois. James Ridgway, London.
- Bowles, M.L., M.D. Hutchison, & J.L. McBride. 1994. Landscape pattern and structure of oak savanna, woodland, and barrens in northeastern Illinois at the time of European settlement. In: Proceedings of the North American Conference on Savannas and Barrens. J. Fralish, ed. Illinois State University.
- Bray, J.R. 1958. The distribution of savanna species in relation to light intensity. *Canadian Journal of Botany* 36:671-681.
- Bray, J.R. 1960. The composition of savanna vegetation in Wisconsin. *Ecology* 41:721-733.
- Brendel, F. 1887. *Flora Peoriana*. J.W. Franks & Sons, Peoria, Illinois.
- Burger, G.V. J.E. Ebinger, & G.S. Wilhelm (eds). 1991. *Proceedings of the Oak Woods Management Workshop*. Eastern Illinois University, Charleston.
- Curtis, J.T. 1959. *The vegetation of Wisconsin - an ordination of plant communities*. The University of Wisconsin Press, Madison.
- Deam, C.C. 1940. *Flora of Indiana*. Indiana Department of Conservation, Department of Forestry. Indianapolis.
- DeSelm, H. 1981. Characterization of some southeastern barrens with special reference to Tennessee. Pages 86-88 in: Proceedings of the Sixth North American Prairie Conference. Ronald Stuckey & K.J. Reese, eds. Ohio Biological Survey Notes No. 15.
- DeSelm, H. 1986. Natural forest openings on uplands of the eastern United States. 1986. Pages 366-375 in: Wilderness and natural areas in the eastern United States: a management challenge. D.L. Kulhavy & D.N. Conner, eds. Stephen F. Austin State University, Nacagdoches, Texas.
- Engelmann, H. 1863. Remarks upon the causes producing the different characters of vegetation known as prairies, flats, and barrens in southern Illinois, with special reference to observations made in Perry and Jackson counties. *American Journal of Science* 86:384-396.
- Faber-Langendoen, D. *in press*. Characterization and classification of oak savanna and woodland communities. In: Proceedings of the 1993 Oak Savanna Conference. F. Stearns, ed. Illinois Environmental Protection Agency, Chicago.
- Farnham, E.W. 1846. *Life in prairie land*. Arno Press, New York (1972).
- Gerhard, F. 1857. *Illinois as it is*. Keen and Lee, Chicago, Illinois.
- Gleason, H.A. 1913. The relation of forest distribution and prairie fires in the middle west. *Torreyana* 13:173-181.
- Gleason, H.A. 1922. The vegetational history of the Middle West. *Annals of the Association of American Geographers* 12:39-85.
- Hall, I.V. & L.E. Aalders. 1968. The botanical composition of two barrens in Nova Scotia. *Naturaliste Canadien* 95:393-396.
- Higley, W.K. & C.S. Raddin. 1891. The flora of Cook County, Illinois, and a part of Lake County, Indiana. *Bulletin of the Chicago Academy of Science* 2:i-xxiii, 1-168.

- Hilgard, E.W. n.d. Botanical features of the prairies of Illinois in ante-railroad days. Typescript at the Illinois Historical Survey, Urbana.
- Homoya, M. 1994. Barrens as an ecological term: an overview of usage in the scientific literature. In: Proceedings of the North American Conference on Savannas and Barrens. J. Fralish, ed. Illinois State University.
- Hus, H. 1908. An ecological cross section of the Mississippi River in the region of St. Louis, Missouri. Nineteenth Annual report of the Missouri Botanical Garden.
- Hutchison, M.D., S. Olson, & T. Vogt. 1986. A survey of the barrens region in Massac and Pope counties, Illinois. Natural Land Institute, Belknap.
- Hutchison, M. 1987. Presettlement barrens in Illinois. Pages 77-90 in: Proceedings of the 8th Northern Illinois Workshop. Forest Preserve District of Illinois. Joliet.
- Keith, J.H. 1980. Presettlement barrens of Harrison and Washington counties, Indiana. Pages 17-25 in: Proceedings of the Seventh North American Prairie conference. C.L. Kucera, ed. Southwest Missouri State University, Springfield.
- Kilburn, P.D. 1959. The forest-prairie ecotone in northeastern Illinois. *American Midland Naturalist* 62:206-217.
- Leach, M.K. 1994. Research begins on restoring oak savanna groundlayer (Wisconsin). *Restoration & Management Notes* 12:83-84.
- Madany, M.H. 1981. A floristic survey of savannas in Illinois. Pages 177-181 in: Proceedings of the Sixth North American Prairie Conference. R.L. Stuckey & K.J. Reese, eds. Ohio Biological Survey Notes 15. Columbus.
- Matlack, G.R., D.J. Gibson, & R.E. Good. Clonal propagation, local disturbance, and the structure of vegetation: ericaceous shrubs in the Pine Barrens of New Jersey. *Biological Conservation* 63:1-8.
- Mead, S.B. 1846. Catalogue of plants growing spontaneously in the State of Illinois, the principal part near Augusta, Hancock County. *Prairie Farmer* 6:35-36, 60, 93, 119-122.
- Mitchell, S.A. 1838. Illinois in 1837 & 1838. S. Augustus Mitchell, and Grigg and Elliot, Philadelphia, Penn.
- Moran, R. 1976. Presettlement vegetation of Lake County, Illinois. Pages 12-18 in: Fifth Midwest Prairie Conference. D.Glenn-Lewin and R. Landers, eds. Iowa State University, Ames.
- Moran, R.C. 1980. Presettlement (1830) vegetation of DeKalb, Kane and DuPage counties, Illinois. M.S. Thesis. Southern Illinois University, Carbondale.
- Packard, S. 1991. Rediscovering the tallgrass savanna. Pages 55-67 in: Proceedings of the Oak Woods Management Workshop. G.V. Burger, J.E Ebinger, and G.S. Wilhelm eds. Eastern Illinois University, Charleston.
- Packard, S. 1993. Restoring oak ecosystems. *Restoration & Management Notes* 11:5-16.
- Peck, J.M. 1834. A gazetteer of Illinois. J.M. Goudy, Jacksonville.
- Ridgway, R. 1873. Notes on the vegetation of the lower Wabash valley. *American Naturalist* 7:154-157.
- Schroeder, W.A. 1982. Presettlement prairie of Missouri. Natural History Series, No.2 Missouri Department of Conservation, Jefferson City.

- Short, C.W. 1845. Observations on the botany of Illinois, more especially in reference to the autumnal flora of the prairies. *Western Journal of Medicine and Surgery*, New Series 3:185-198..
- Strang, R.M. 1972. Ecology and land use of the barrens of western Nova Scotia. *Canadian Journal of Forest Research* 2(3):276-290.
- Stearns, F.W. (ed.) *in press*. Proceedings of the 1993 Oak Savanna Conference. Illinois Environmental Protection Agency, Chicago.
- Stout, A.B. 1946. The bur oak openings in southern Wisconsin. *Transactions of the Wisconsin Academy of Science, Arts, and Letters* 36:141-161.
- Vestal, A.G. 1936. Barrens vegetation in Illinois. *Transactions of the Illinois Academy of Science* 29:29-80.
- White, J. & M. Madany. 1978. Classification of natural communities in Illinois. Appendix 30 in: Illinois Natural areas inventory technical report. Department of Landscape Architecture, University of Illinois, Urbana-Champaign & Natural Land Institute, Rockford, Illinois.
- White, J. 1994. How the terms *savanna*, *barrens*, and *oak openings* were used in early Illinois. In: Proceedings of the North American Conference on Savannas and Barrens. J. Fralish, ed. Illinois State University.
- Williams, D.L. 1981. Reconstruction of prairie peninsula vegetation and its characteristics from descriptions before 1860. Pages 83-86 in: Proceedings of the Sixth North American Prairie Conference. Ronald Stuckey & K.J. Reese, eds. Ohio Biological Survey Notes No. 15.
- Woods, J. 1822. Two years residence on the English Prairie of Illinois. Edited by P.M. Angle and published in 1968 by The Lakeside Press, Chicago.

AUTHOR AFFILIATION

Marlin Bowles and Jeanette McBride are with The Morton Arboretum, Lisle, IL




Figure 1. Conceptual landscape model for savanna and barrens development on glaciated silt-loam soils. Horizontal axis: eastward-moving landscape prairie fires cause degeneration of forest (*sensu* Gleason 1922) into barrens or savanna, and are regulated by climate and landscape features. Cool moist conditions reduce fire effects and barrens and savanna encroach on prairie, warm dry conditions increase fire effects and allow barrens and savanna to encroach on forest; firebreaks such as rugged topography or watercourses would affect the landscape pattern of this vegetation. Vertical axis: landscape processes (*e.g.* fire frequency or severity and degree of fire protection) or biological characteristics (*e.g.* fire resistance, dispersal, and colonization rates of different tree and shrub species) favor either woody understory ("Barrens Pathway") or overstory ("Savanna Pathway"). The different pathways may co-occur or integrate. Grub barrens are derived from direct degeneration of forest, or from secondary degeneration of savanna.

APPENDIX I. Historic barrens plant species checklist. Sources: species associated with "barrens" in Mead (1846), "copse" in Brendel (1887), "copse" or "thicket" in Higley & Raddin (1891) and "thicket" in Hus (1908).

<i>Acalypha virginica</i>	<i>Danthonia spicata</i>	<i>Liparis lilifolia</i>	<i>Rhus glabra</i>
<i>Agastache nepetoides</i>	<i>Delphinium carolinianum</i>	<i>Lonicera prolifera</i>	<i>Ribes missouriense</i>
<i>Agastache scrophulariaefolia</i>	<i>Delphinium tricorne</i>	<i>Lycopus americanus</i>	<i>Rosa blanda</i>
<i>Agrimonia parvula</i>	<i>Desmodium canadense</i>	<i>Lysimachia ciliata</i>	<i>Rosa carolina</i>
<i>Agrimonia pubescens</i>	<i>Desmodium canescens</i>	<i>Malus coronaria</i>	<i>Rosa setigera</i>
<i>Allium tricoccum</i>	<i>Desmodium cuspidatum</i>	<i>Melica niens</i>	<i>Rosa sulfata</i>
<i>Ampelamus albidus</i>	<i>Desmodium glutinosum</i>	<i>Menispermum canadense</i>	<i>Rubus allegheniensis</i>
<i>Amphicarpa bracteata</i>	<i>Desmodium illinoense</i>	<i>Monarda fistulosa</i>	<i>Rubus flagellaris</i>
<i>Amphicarpa comosa</i>	<i>Desmodium paniculatum</i>	<i>Napaea dioica</i>	<i>Rubus occidentalis</i>
<i>Andropogon furcatus</i>	<i>Desmodium sessilifolium</i>	<i>Onoclea sensibilis</i>	<i>Sabatia angularis</i>
<i>Andropogon scoparius</i>	<i>Dioscorea villosa</i>	<i>Orobanche uniflora</i>	<i>Sambucus canadensis</i>
<i>Anemone cylindracea</i>	<i>Dodecatheon meadii</i>	<i>Ostrya virginiana</i>	<i>Sanguinaria canadensis</i>
<i>Apocynum androsaemifolium</i>	<i>Echinacea purpurea</i>	<i>Oxalis cymosa</i>	<i>Sanicula marilandica</i>
<i>Apocynum cannabinum</i>	<i>Elymus canadensis</i>	<i>Oxalis stricta</i>	<i>Salix discolor</i>
<i>Aquilegia canadensis</i>	<i>Elymus virginicus</i>	<i>Oxalis violacea</i>	<i>Salix rigida</i>
<i>Arisaema dracontium</i>	<i>Epilobium angustifolium</i>	<i>Panicum clandestinum</i>	<i>Scrophularia marilandica</i>
<i>Arisaema triphyllum</i>	<i>Erigeron annuus</i>	<i>Panicum depauperatum</i>	<i>Scutellaria ovata</i>
<i>Asclepias exaltata</i>	<i>Erigeron pulchellus</i>	<i>Panicum dichotomum</i>	<i>Scutellaria parvula</i>
<i>Asclepias purpurascens</i>	<i>Erythronium albidum</i>	<i>Panicum latifolium</i>	<i>Silene stellata</i>
<i>Asclepias tuberosa</i>	<i>Euonymus atropurpureus</i>	<i>Panicum scribnerianum</i>	<i>Silphium terebinthinaceum</i>
<i>Asclepias verticillata</i>	<i>Eupatorium altissimum</i>	<i>Paronychia canadensis</i>	<i>Silphium integrifolium</i>
<i>Aster azureus</i>	<i>Eupatorium purpureum</i>	<i>Parthenium integrifolium</i>	<i>Silphium perfoliatum</i>
<i>Aster drummondii</i>	<i>Eupatorium rugosum</i>	<i>Passiflora lutea</i>	<i>Smilacina racemosa</i>
<i>Aster lateriflorus</i>	<i>Eupatorium sessilifolium</i>	<i>Petalostemum candidum</i>	<i>Smilax ecirrhata</i>
<i>Aster novae-angliae</i>	<i>Galium aparine</i>	<i>Petalostemum purpureum</i>	<i>Smilax herbacea</i>
<i>Aster praealtus</i>	<i>Galium circaezans</i>	<i>Phaseolus polystachyus</i>	<i>Smilax hispida</i>
<i>Aster turbinellus</i>	<i>Galium concinnum</i>	<i>Phlox divaricata</i>	<i>Solidago buckleyi</i>
<i>Astragalus canadensis</i>	<i>Gentiana flavida</i>	<i>Phryma leptotachya</i>	<i>Solidago caesia</i>
<i>Aureolaria pedicularia</i>	<i>Gentiana quinquefolia</i>	<i>Physalis subglabrata</i>	<i>Solidago canadensis</i>
<i>Boehmeria cylindrica</i>	<i>Geranium carolinianum</i>	<i>Podophyllum peltatum</i>	<i>Solidago gigantea</i>
<i>Baptisia leucantha</i>	<i>Geranium maculatum</i>	<i>Polygala ambigua</i>	<i>Solidago juncea</i>
<i>Baptisia leucophaea</i>	<i>Gerardia grandiflora</i>	<i>Polygala incarnata</i>	<i>Solidago speciosa</i>
<i>Campanula americana</i>	<i>Gerardia tenuifolia</i>	<i>Polygala polygama</i>	<i>Solidago ulmifolia</i>
<i>Carex blanda</i>	<i>Geum canadense</i>	<i>Polygala sanguinea</i>	<i>Staphylea trifolia</i>
<i>Carex cephalophora</i>	<i>Geum laciniatum</i>	<i>Polygala senega</i>	<i>Strophostyles helvola</i>
<i>Carex jamesii</i>	<i>Hackelia virginiana</i>	<i>Polygala verticillata</i>	<i>Symphoricarpos orbiculatus</i>
<i>Carex richardsonii</i>	<i>Hamamelis virginiana</i>	<i>Polygonum dumetorum</i>	<i>Syntherisma bullii</i>
<i>Carya cordiformis</i>	<i>Helianthus decapetalus</i>	<i>Polygonum scandens</i>	<i>Taenidia integerrima</i>
<i>Carya glabra</i>	<i>Helianthus divaricatus</i>	<i>Polygonum virginianum</i>	<i>Tephrosia virginiana</i>
<i>Cassia fasciculata</i>	<i>Helianthus giganteus</i>	<i>Polytaenia nuttallii</i>	<i>Thalictrum dasycarpum</i>
<i>Castilleja coccinea</i>	<i>Helianthus grosseserratus</i>	<i>Populus tremuloides</i>	<i>Tradescantia virginica</i>
<i>Ceanothus americanus</i>	<i>Helianthus hirsutus</i>	<i>Potentilla arguta</i>	<i>Trillium recurvatum</i>
<i>Celastrus scandens</i>	<i>Helianthus occidentalis</i>	<i>Potentilla simplex</i>	<i>Trillium sessile</i>
<i>Cercis canadensis</i>	<i>Helianthus strumosus</i>	<i>Prenanthes alba</i>	<i>Triosteum perfoliatum</i>
<i>Chaerophyllum procumbens</i>	<i>Heliopsis helianthoides</i>	<i>Prunella vulgaris</i>	<i>Veratrum woodii</i>
<i>Chamaesyce maculata</i>	<i>Heracleum lanatum</i>	<i>Prunus americana</i>	<i>Verbesina alternifolia</i>
<i>Cirsium altissimum</i>	<i>Humulus americanus</i>	<i>Prunus serotina</i>	<i>Verbesina helianthoides</i>
<i>Cirsium discolor</i>	<i>Hydrastis canadensis</i>	<i>Prunus virginiana</i>	<i>Vernonia missurica</i>
<i>Claytonia virginica</i>	<i>Hypericum punctatum</i>	<i>Psoralea onobrychis</i>	<i>Viburnum lentago</i>
<i>Comandra richardsiana</i>	<i>Hypericum sphaerocarpum</i>	<i>Psoralea tenuiflora</i>	<i>Viburnum prunifolium</i>
<i>Commelina erecta</i>	<i>Ipomoea pandurata</i>	<i>Pteridium aquilinum</i>	<i>Vicia americana</i>
<i>Convolvulus sepium</i>	<i>Isanthus brachiatus</i>	<i>Pycnanthemum flexuosum</i>	<i>Viola cuculata</i>
<i>Convolvulus spithameus</i>	<i>Lactuca canadensis</i>	<i>Pycnanthemum pilosum</i>	<i>Viola palmata</i>
<i>Coreopsis palmata</i>	<i>Lactuca ludoviciana</i>	<i>Pycnanthemum virginianum</i>	<i>Viola pedata</i>
<i>Coreopsis tripteris</i>	<i>Lathyrus palustris</i>	<i>Quercus bicolor</i>	<i>Viola pedatifida</i>
<i>Cornus alternifolia</i>	<i>Lathyrus venosus</i>	<i>Quercus imbricaria</i>	<i>Viola sagittata</i>
<i>Cornus drummondii</i>	<i>Lechea minor</i>	<i>Quercus marilandica</i>	<i>Vitis vulpina</i>
<i>Cornus obliqua</i>	<i>Lespedeza capitata</i>	<i>Quercus velutina</i>	<i>Zanthoxylum americanum</i>
<i>Cornus rugosa</i>	<i>Lespedeza procumbens</i>	<i>Ranunculus abortivus</i>	
<i>Corylus americana</i>	<i>Lespedeza violacea</i>	<i>Ranunculus recurvatus</i>	
<i>Cynoglossum virginianum</i>	<i>Lespedeza virginica</i>	<i>Rhamnus lanceolata</i>	
<i>Cypripedium calceolus</i>	<i>Liatris scariosa</i>	<i>Rhus aromatica</i>	
<i>Cypripedium reginae</i>	<i>Lilium michiganense</i>	<i>Rhus copallina</i>	