

Mean (+std. err.) dominance (relative basal area) for canopy species sampled in 1996 in maple, red oak, and white oak stands along transect lines established by the INAI. Species ranked by overall stand means.

Species	Maple stands		Red oak stands		White oak stands	
	Avg.	Std err	Avg.	Std err	Avg.	Std err
<i>Quercus alba</i>	7.437	2.184	24.947	3.690	47.188	7.292
<i>Quercus rubra</i>	15.696	4.193	41.340	6.221	16.761	5.253
<i>Acer saccharum</i>	45.071	4.818	5.282	2.928	3.842	2.495
<i>Fraxinus americana</i>	7.980	2.124	8.052	1.411	3.517	2.054
<i>Tilia americana</i>	10.117	2.165	3.159	0.980	0.871	0.747
<i>Ulmus rubra</i>	3.810	1.292	4.704	1.360	3.779	2.296
<i>Quercus velutina</i>	0.097	0.100	1.972	1.151	5.223	3.359
<i>Prunus serotina</i>	0.288	0.102	1.893	0.756	4.240	2.104
<i>Ostrya virginiana</i>	2.884	1.669	1.443	0.406	1.930	1.648
<i>Quercus macrocarpa</i>	1.292	0.652	1.376	1.076	2.739	2.144
<i>Quercus bicolor/mac.</i>	0.985	0.759	2.193	1.822	1.840	1.089
<i>Juglans nigra</i>	1.913	1.107	0.764	0.334	1.885	1.071
<i>Carya ovata</i>	0.198	0.102	0.898	0.730	2.046	0.821
<i>Ulmus americana</i>	0.473	0.473	0.735	0.735	1.307	0.957
<i>Quercus coccinea</i>	0.100	0.100	0.071	0.071	1.503	1.064
<i>Carya cordiformis</i>	0.113	0.113	0.414	0.187	0.695	0.695
<i>Populus deltoides</i>	0.539	0.408	0.265	0.190	0.000	0.000
<i>Fraxinus quadrangulata</i>	0.242	0.242	0.000	0.000	0.420	0.420
<i>Juglans cinerea</i>	0.100	0.100	0.433	0.218	0.000	0.000
<i>Betula papyrifera</i>	0.333	0.333	0.000	0.000	0.000	0.000
<i>Populus grandidentata</i>	0.200	0.200	0.000	0.000	0.100	0.100
<i>Fraxinus pennsylvanica</i>	0.100	0.073	0.000	0.000	0.078	0.078
<i>Carpinus caroliniana</i>	0.118	0.118	0.100	0.100	0.000	0.000
<i>Acer rubrum</i>	0.000	0.000	0.000	0.000	0.092	0.092
<i>Prunus virginiana</i>	0.068	0.068	0.000	0.000	0.000	0.000
<i>Crataegus pruinosa</i>	0.000	0.000	0.100	0.100	0.000	0.000
<b>Mean stand basal area</b>	<b>25.390</b>	<b>1.700</b>	<b>24.340</b>	<b>1.420</b>	<b>25.350</b>	<b>0.980</b>